

Stop Your Back Pain



Benefits

1 80+% Improvement

2 FDA-approved 2010

3 Outpatient Surgery



LESS™ Exposure Spine Surgery

< 3 cm incision

LESS™ is More

Interspinous Fixation For **Spinal Stenosis & Fusion**

CPT Code 22612, 22899, 63047



LESS™ Exposure Advanced Spine Technologies (LEAST)



Spine Pain

Symptoms

- Back, buttock or leg pain
- Difficulty walking
- Numbness in your legs, calves, or buttocks
- Weakness and / or loss of balance

Potential Causes

- Degenerative disc
- Spinal Stenosis
- Herniated Disc
 - Spondylolisthesis
 - Scoliosis
 Spinal Fractures
 - Painful Facet Joints



Indications For Surgery



- Radicular pain or neurogenic claudication resulting in functional impairment
 - Failed 3 months of conservative care (NSAIDS, Physical therapy, Activity modications, Injections).
 - Lumbar spinal stenosis on MRI/CT
 - X-Ray/CT/MRI evidence of instability

Ways To Use InSpan







Post-Op Information

- Follow up with your doctor
- Keep incision dry for 7-10 days
- Take antibiotics for 5-7 days
- Pain medications as prescribed
- NO NSAIDS
- NO smoking

- Walk as tolerated
- No lifting above 10 lbs
- Physical therapy
- · Use brace as prescribed
- Consult your doctor: bending, driving, exercising





" My back pain was unbearable and I just could not stand, InSpan has given me back my life " - Patient

Not every patient will qualify for InSpan or have the best results



www.NANISX.com www.MyInSpan.com

