


## Stop Sacroiliac Joint Pain

**LESS  
Recovery Time**

*Get Back  
To Action*



## Benefits

- 1 98+% Success Rate**
- 2 FDA-approved 2015**
- 3 Outpatient Surgery**



Nanofuse used with Sacrix  
**NANOFUSE**  
BIOLOGICS  
Synthetic Bioactive Grafts  
[www.Nanofusebiologics.com](http://www.Nanofusebiologics.com)

**LESS<sup>™</sup> Exposure Spine Surgery**

**LESS<sup>™</sup> is More**

**Percutaneous  
Lateral-Oblique Transfixation**

CPT Code 27279



# Diagnosis

## Symptoms

- Back, buttock or leg pain
- Difficulty walking
- Numbness in your legs, calves, or buttocks
- Weakness and / or loss of balance

## Potential Causes

- Degenerative disc
- Painful facet joints
- Spinal stenosis
- Spondylolisthesis
- Scoliosis
- Spinal Fractures
- Failed prior fusion



## Provocative Tests

COMPRESSION TEST



FABER TEST



GAENSLER'S TEST




THIGH THRUST TEST



DISTRACTION TEST



## Indications For Surgery



- ◆ Pain longer than 4-6 months
- ◆ At least 3 positive provocative tests
- ◆ X-Ray & CT/MRI imaging
- ◆ 75% improvement with injection
- ◆ Steroid injection at least once
- ◆ Pain over the SI Joint

## Post-Op

- Follow up with your doctor
- Keep incision dry for 7-10 days
- Take antibiotics for 5-7 days
- Pain medications as prescribed
- NO NSAIDS
- NO smoking
- Walk as tolerated
- No lifting above 10 lbs
- Physical therapy
- Use brace as prescribed
- Consult your doctor: bending, driving, exercising



***"Since having the Sacrix procedure, I am able to be active and sleep at night without pain medication"***

- Patient

Not every patient will qualify for Sacrix or have the best results

